

Curves of constant diameter and inscribed polygons

Mathieu Baillif


Let Γ be a simple closed curve in the Euclidean plane. Say that a polygon S is *inscribed in Γ at x* if all the vertices of S lie on Γ and one is x . A line segment is here considered as a 2-gon. We say that Γ has property $(C_n(D))$ (for some $n \geq 2$, $D > 0$) if

$$\forall x \in \Gamma \text{ there is a unique regular } n\text{-gon with edges length } D \text{ inscribed in } \Gamma \text{ at } x. \quad (C_n(D))$$

Notice that $(C_2(D))$ is equivalent to the following since Γ is simple and closed ($\|\cdot\|$ is the Euclidean norm):

$$\forall x \in \Gamma \quad \exists! y(x) \in \Gamma \text{ with } \|x - y(x)\| = D, \text{ and} \quad (C(D)) \\ \text{if } z \neq y(x), \|x - z\| < D.$$

If one drops the unicity assumption, $(C(D))$ is the property of having *constant diameter*, which is in fact equivalent (for closed curves in the plane) to having *constant width* or *constant breadth* (for the definitions and the proof of the equivalence, see [8, chap. 25]). It is a surprise to many (including me !) that curves of constant diameter other than the circle do exist. The simplest examples (due to Reuleaux [9]) are pictured below. They are built with circle arcs whose centers are marked with a black dot. Notice that these curves are not $\mathcal{C}^2$.



The theory of curves of constant breadth has generated a considerable literature, starting with Euler [3] and including works by Hurwitz [4], Minkowski [6], Blaschke (five articles in [1]) and many others. These curves have many interesting properties, the most startling being perhaps that their perimeter is π times their diameter. See [8, chap. 25] for an elementary account, or [5] for a more thorough, though old, presentation. For $n \geq 3$, to

our knowledge the property $(C_n(D))$ has not been investigated (but see [5, p. 61], and [2] for a related problem).

In this note, we shall prove the theorem below, using only basic differential geometry (tangent, curvature, and so on). We recall that a curve is *regular* if it is $\mathcal{C}^1$ with non vanishing derivative.

THEOREM. *i) For all $n \geq 2$ and $D > 0$, there are $\mathcal{C}^\infty$ regular simple closed non circular curves Γ with property $(C_n(D))$.*

ii) The circle of radius $D/2$ is the only $\mathcal{C}^2$ regular simple closed curve which satisfies both $(C_4(\frac{D}{\sqrt{2}}))$ and $(C(D))$.

REMARKS. For i) $n = 2$, such $\mathcal{C}^\infty$ curves abound in the literature, see for instance [10] or [7] for one given by a polynomial equation. We however give a short self contained proof that yields simple explicit examples. For ii), notice that a square with edges length $D/\sqrt{2}$ has diagonal D . This result gives a definition of the circle involving only Euclidean distance between points *on* the circle, while most usual definitions refer to points *off* the circle.

Proof of i) for $n = 2$. We shall prove the following:

PROPOSITION 1. *Let $G : \mathbb{R} \rightarrow \mathbb{R}^2$ be $\mathcal{C}^2$ such that*

a) $G(\theta + \pi) = G(\theta)$,

b) $G'(\theta) = r(\theta) \cdot (-\sin \theta, \cos \theta)$, where $|r(\theta)| < \frac{D}{2}$.


Then, the curve $\gamma(\theta) = G(\theta) + \frac{D}{2}(\cos \theta, \sin \theta)$ has property $(C(D))$.

To obtain explicit examples, we can take $r(\theta) = a \cdot \sin((2k+1)\theta)$, with $k \geq 1$, $a < D/2$. Integrating, we get

$$G(\theta) = \frac{a}{4} \left(-\frac{\sin(2k\theta)}{k} + \frac{\sin(2(k+1)\theta)}{k+1}, -\frac{\cos(2k\theta)}{k} - \frac{\cos(2(k+1)\theta)}{k+1} \right)$$

which satisfies $G(\theta) = G(\theta + \pi)$. By Proposition 1, the curve $\gamma(\theta) = G(\theta) + \frac{D}{2}(\cos \theta, \sin \theta)$ has property $(C(D))$. One could also take any linear combination of $\cos((2k+1)\theta)$ and $\sin((2k+1)\theta)$ ($k \geq 1$), with small enough coefficients, for $r(\theta)$. Below are pictured the curves of diameter 1 (and G) for

$$r(\theta) = \sin(3\theta)/3 + \cos(3\theta)/5, \sin(5\theta)/2.01, \sin(3\theta)/10 + \cos(7\theta)/2.501.$$




Proof of Proposition 1. By construction, $\gamma(\theta + \pi) - \gamma(\theta)$ is collinear with $(\cos \theta, \sin \theta)$. Thus, $\gamma'(\theta) = (r(\theta) + \frac{D}{2})(-\sin \theta, \cos \theta)$ is normal to $\gamma(\theta + \pi) - \gamma(\theta)$. Moreover, $r(\theta) > -\frac{D}{2}$, $\gamma'(\theta) \neq 0$, thus γ is regular. We could end the proof here by invoking the classical result that a curve such that every normal is a binormal is of constant diameter (see for instance [5]), but let us give a self contained argument. Since $\gamma''(\theta) = r'(\theta) \cdot (-\sin \theta, \cos \theta) - (r(\theta) + \frac{1}{2})(\cos \theta, \sin \theta)$, the curvature of γ is

$$\rho(\theta) = \frac{\|\gamma'(\theta) \wedge \gamma''(\theta)\|}{\|\gamma'(\theta)\|^3} = \frac{1}{|r(\theta) + D/2|} > \frac{1}{D},$$

since $r(\theta) < D/2$. To summarize, γ satisfies:

- 1) $\|\gamma(\theta) - \gamma(\theta + \pi)\| = D$,
- 2) $\gamma'(\theta)$ is normal to $\gamma(\theta) - \gamma(\theta + \pi)$,
- 3) the curvature is $> 1/D$.

We now show that γ satisfies $(C(D))$. If $\epsilon > 0$ is small enough, since the curvature of γ at $\theta + \pi$ is strictly bigger than the one of the circle of radius D and center $\gamma(\theta)$ while their derivatives are collinear, $\gamma(\theta + \pi \pm \epsilon)$ is in the interior of this circle, and thus $\|\gamma(\theta + \pi \pm \epsilon) - \gamma(\theta)\| < \|\gamma(\theta + \pi) - \gamma(\theta)\|$ (see the figure below, on the left). If one “goes away” from $\theta + \pi$, the distance will decrease more and more. Indeed, by construction the segments between $\gamma(\phi)$ and $\gamma(\phi + \pi)$ (for different ϕ) all cross each other. We are thus in the case of the figure below, on the right.




The circle of radius D and center $\gamma(\phi + \pi)$ is dashed. For the same reason as above, the curve will stay (locally) inside this circle. Thus,

$$\|\gamma(\phi) - \gamma(\theta)\| > \|\gamma(\phi + \epsilon) - \gamma(\theta)\|$$

(see figure). The distance decreases thus strictly when ϕ gets away from $\theta + \pi$. $\square$

Proof of i) for $n \geq 3$.


Let $\gamma(\theta) = G(\theta) + R(\cos \theta, \sin \theta)$, where R is the radius of the n -gon with edges length D , and G satisfies $G(\theta + 2\pi/n) = G(\theta)$. By construction, $\{\gamma(\theta + \frac{2k\pi}{n})\}_{k=0 \dots n-1}$ are the vertices of a regular n -gon with edges length D (see the figure on the right). We will show that if G and its derivative are small enough, there are no other such n -gon inscribed in Γ at $\gamma(\theta)$.



For this, write $f_\theta(\phi) = \|\gamma(\theta) - \gamma(\theta + \phi)\|$ and suppose first that γ is a circle (i.e. that G is constant). We fix $\epsilon > 0$ such that if $\phi \in [-\epsilon, \epsilon]$ then $f_\theta(\phi) < D$ and if $\phi \in [\pi - \epsilon, \pi + \epsilon]$, $f_\theta(\phi) > D$. This property will remain true (for all θ and for the same ϵ) if G is small enough. If γ is a circle, $f_\theta(\phi)$ is strictly increasing on $]0, \pi[$ and strictly decreasing on $]\pi, 2\pi[$. Thus, if G and its derivative are small, f_θ will be strictly increasing on $]\epsilon, \pi - \epsilon[$ and strictly decreasing on $]\pi + \epsilon, 2\pi - \epsilon[$ for all θ . There are therefore only two points at distance D from $\gamma(\theta)$, so there is a unique n -gon with edges length D inscribed in the curve at $\gamma(\theta)$ (if G and its derivative are small enough). $\square$

Proof of ii). Let Γ be a $\mathcal{C}^2$ regular simple closed curve satisfying $(C_4(D/\sqrt{2}))$ and $(C(D))$. We take $D = 1$ for simplicity. Let γ parametrise Γ by arc length counterclockwise.

If $x \in \Gamma$, denote by $y(x)$ the unique point of Γ for which $\|x - y(x)\| = 1$. If $\gamma(t) = x$, $\gamma'(t)$ must be normal to $x - y(x)$, because the curve must be totally inside the “dashed eye” of the figure on the right, and in fact, $y(x)$ is the point on the normal at x at distance 1 from x . Moreover, the curvature must be > 1 .




Since the unit normal vector at $\gamma(t)$ is $n(t) = (-\gamma'_2(t), \gamma'_1(t))$, $y(\gamma(t)) = \gamma(t) + n(t)$ is differentiable in t . (Notice that we cannot use the implicit function theorem here, because the derivative of $\|x - z\|$ vanishes precisely at $z = y(x)$.) Moreover, the angle between the oriented line segment $[\gamma(t), y(\gamma(t))]$ and the horizontal axis strictly increases. Given $\theta \in [0, 2\pi]$, there is thus a unique (oriented) line segment $[x, y(x)]$ making an angle θ with the horizontal axis, we define $G(\theta)$ as its middle point. Since $y(x)$ is differentiable with respect to x , $G(\theta)$ is differentiable with respect to θ . Then, $\tilde{\gamma}(\theta) = G(\theta) + \frac{1}{2}(\cos \theta, \sin \theta)$ is a parametrisation of Γ (see the figure below, on the left). By definition, $y(\tilde{\gamma}(\theta)) = \tilde{\gamma}(\theta + \pi)$, and since the tangent of Γ at x is normal to $x - y(x)$, we have

$$\langle \tilde{\gamma}'(\theta) | (\cos \theta, \sin \theta) \rangle = \langle G'(\theta) | (\cos \theta, \sin \theta) \rangle = 0,$$

which implies $G'(\theta) = r(\theta)(-\sin \theta, \cos \theta)$ for some continuous r .

Now, since Γ has property $(C_4(1/\sqrt{2}))$, there is a unique square $S(\theta)$ with edges length $\frac{1}{\sqrt{2}}$ inscribed in Γ at $\tilde{\gamma}(\theta)$. Since $\tilde{\gamma}(\theta + \pi)$ is the unique point of Γ at distance 1 from $\tilde{\gamma}(\theta)$, $\tilde{\gamma}(\theta + \pi)$ is the vertex of $S(\theta)$ diagonal to $\tilde{\gamma}(\theta)$ ($S(\theta)$ has diagonal 1). Thus, $G(\theta)$ is also the center of $S(\theta)$, which implies $G(\theta + \pi/2) = G(\theta)$ (see figure below, on the right).



Therefore, $G'(\theta+\pi/2) = G'(\theta)$, i.e. $r(\theta+\pi/2)(\cos \theta, \sin \theta) = r(\theta)(-\sin \theta, \cos \theta)$, so $r(\theta) = 0$, $G(\theta)$ is constant, and hence Γ is a circle. $\square$

This proof can be easily generalized to show that if a $\mathcal{C}^2$ regular closed simple curve has both properties $(C_{2n}(\tilde{D}))$ and $(C(D))$, with D two times the radius of the regular $2n$ -gon with edges length $\tilde{D}$, then it is the circle. This however leaves open the following:

QUESTIONS. Are there $D, \tilde{D} > 0$ and curves other than the circle which have property $(C(D))$ and $(C_n(\tilde{D}))$ for some $n \geq 3$? Are there (non regular or non $\mathcal{C}^2$) curves with properties $(C(D))$ and $(C_4(\frac{D}{\sqrt{2}}))$?

We finish with the following proposition, which gives a motivation for the “!” in the definition of $(C(D))$:

PROPOSITION 2. *Let Γ be a continuous closed piecewise regular curve satisfying $(C(D))$. Then, Γ is regular.*

Piecewise regular means that the curve is regular everywhere, except possibly at a finite number of points. The proof is left as an exercise. (*Hint:* Show that $y(x)$ is continuous in x , then that Γ cannot have corners.) Notice that taking $b = 0$ in the first figure of this paper yields a curve of constant diameter with corners (the well known Reuleaux triangle).

ACKNOWLEDGEMENTS. I wish to thank D. Cimasoni, G. Wanner, and especially B. Dudez, librarian at the math department in Geneva, for his bibliographical help.

References

- [1] W. Blaschke. *Gesammelte Werke*, volume 3. Thales-Verl., Essen, 1982–1986.
- [2] H. G. Eggleston and S. J. Taylor. On the size of equilateral triangles which may be inscribed in curves of constant width. *J. London Math. Soc.*, 27:438–448, 1952.
- [3] L. Euler. De curvis triangularibus. *Acta Acad. Petropol.* 1778, II:3–30, 1780. Reprinted in *Opera Omnia*, vol. 28.
- [4] A. Hurwitz. Sur quelques applications géométriques des séries de Fourier. *Ann. Sci. ENS*, pages 357–408, 1902. tome 19.
- [5] Ch. Jordan and R. Fiedler. *Contribution à l'étude des courbes convexes fermées*. Hermann, Paris, 1912.
- [6] H. Minkowski. Über die Körper konstanter Breite. In *Gesammelte Abhandlungen*, volume II, pages 277–279. Chelsea Publ., New York, 1967. Reprint of the Leipzig Edition, 1911.

- [7] S. Rabinowitz. A polynomial curve of constant width. *Missouri J. Math.*, 9(1):23–27, 1997.
- [8] H. Rademacher and O. Toeplitz. *The enjoyment of mathematics*. Princeton University Press, Princeton, N.J., 1957.
- [9] F. Reuleaux. *Theoritishe Kinematik*. Braunschweig, 1875. Available at http://historical.library.cornell.edu/kmoddl/toc_reuleaux1.html.
- [10] S. A. Robertson. Smooth curves of constant width and transnormality. *Bull. London math. soc.*, 16(3):264–274, 1984.